

**Association between
categorical variables
(Percentages)
Chi square (χ^2) test**

Dr Abubakir Majeed
Assist. Prof. of Community Medicine

Outline

- 1) Construct 2-way table to examine association between two categorical variables.
- 2) Conduct Chi Square (χ^2) test to assess evidence for association between two or more categorical variables.

Constructing a two-way table

- Shows distribution of (relationship between) 2 categorical variables.
- Example: Relationship between physical exercise and the sex of individuals?
- If rows are independent variable, use row %'s.
- **2x2 table**

Sex	Exercise		No exercise		Total	
	No.	%	No.	%	No.	%
Male	31	75.6	10	24.4	41	100
Female	101	83.5	20	16.5	121	100
Total	132	81.5	30	18.5	162	100

Another example

- Drug A: Of 93 patients, 49 had response
- Drug B: Of 91 patients, 18 had response
- Construct a two-way table
- **2x2 table**

Drug	Tumor response				Total	
	Yes		No		No.	%
	No.	%	No.	%		
Drug A						
Drug B						
Total						

Another example

- Drug A: out of 93 patients, 49 had response
- Drug B: Out of 91 patients, 18 had response
- Construct a two-way table
- **2x2 table**

Drug	Tumor response				Total	
	Yes		No		No.	%
	No.	%	No.	%		
Drug A	49				93	
Drug B	18				91	
Total	67				184	

Another example

- Drug A: out of 93 patients, 49 had response
- Drug B: Out of 91 patients, 18 had response
- Construct a two-way table
- **2x2 table**

Drug	Tumor response				Total	
	Yes		No		No.	%
	No.	%	No.	%		
Drug A	49		44		93	
Drug B	18		73		91	
Total	67		117		184	

Another example

- Drug A: out of 93 patients, 49 had response
- Drug B: Out of 91 patients, 18 had response
- Construct a two-way table
- **2x2 table**

Drug	Tumor response				Total	
	Yes		No		No.	(%)
	No.	(%)	No.	(%)		
Drug A	49	(53%)	44	(47%)	93	(100%)
Drug B	18	(20%)	73	(80%)	91	(100%)
Total	67	(36%)	117	(64%)	184	(100%)

Larger tables

- **3x3 table**

Age group	Fever after operation						Total	
	Mild		Moderate		Severe		No.	%
	No.	%	No.	%	No.	%		
<30 Y	37	59	14	22	12	19	63	100
30-45 Y	18	33	17	31	19	35	54	100
>45 Y	24	50	14	29	10	21	48	100
Total	79		45		41		165	

Association between two variables

- Two variables are associated if distribution of one varies according to value of other
- Knowing value of one variable tells us something about value of other
- In example,
Knowing sex of student will tell us something about physical exercise (association).

- Usually examine distribution of dependent variable according to levels of independent variable
- Distribution of physical exercise (dependent) across sex (independent)

Sex	Exercise		No exercise		Total	
	No.	%	No.	%	No.	%
Male	31	75.6	10	24.4	41	100
Female	101	83.5	20	16.5	121	100
Total	132	81.5	30	18.5	162	100

- Distribution of physical exercise differs according to sex but.....by more than we expect by chance??.....

Example: Gender and Exercise among students

Sex	Exercise		No exercise		Total	
	No.	%	No.	%	No.	%
Male	31	75.6	10	24.4	41	100
Female	101	83.5	20	16.5	121	100
Total	132	81.5	30	18.5	162	100

75.6% of male students exercise regularly

83.5% of female students exercise regularly

Is there a real difference or it is due to chance?

If no association, H_0 :

Sex	Exercise		No exercise		Total	
	No.	%	No.	%	No.	%
Male		?		?	41	100
Female		?		?	121	100
Total	132	81.5	30	18.5	162	100

If no association, H_0 :

Sex	Exercise		No exercise		Total	
	No.	%	No.	%	No.	%
Male		81.5		18.5	41	100
Female		81.5		18.5	121	100
Total	132	81.5	30	18.5	162	100

Significance test for association

- Examining percentages indicates whether association may exist between exposure and disease
- But is association likely to be real or due to sampling variability?
- Need a

Significance test for association

- Examining percentages indicates whether association may exist between exposure and disease
- But is association likely to be real or due to sampling variability?
- Need a **significance test.**
- **Null hypothesis (H_0): “no association between the two variables”**
- **H_0 : distribution of physical exercise is same in each group (male and female).**

Significance test for comparing proportions

- The test is called Chi Square (χ^2) test
- **Step 1 – Calculate expected table**
For H_0 , as there is no real association
 - Difficult way
 - Quick way
- **Step 2 – Calculate χ^2**
- **Step 3 – Obtain p-value and interpret it**

Note: Steps 1 & 2 can be done in one quick step only for 2x2 tables

Step 1 – Calculate expected table

Sex	Exercise		No exercise		Total	
	No.	%	No.	%	No.	%
Male		81.5		18.5	41	100
Female		81.5		18.5	121	100
Total	132	81.5	30	18.5	162	100

132/162 (81.5%) exercise

30/162 (18.5%) do not exercise

Under H_0 expect same percentage for both sexes

Sex	Exercise		No exercise		Total	
	No.	%	No.	%	No.	%
Male		81.5		18.5	41	100
Female		81.5		18.5	121	100
Total	132	81.5	30	18.5	162	100

Difficult way

Male

Expect 81.5% of male to exercise (81.5% of $41=81.5 \times 41/100=33.4$)

Expect 18.5% of male to not exercise (18.5% of $41=18.5 \times 41/100=7.6$)

Female

Expect 81.5% of female to exercise (81.5% of $121=81.5 \times 121/100=98.6$)

Expect 18.5% of male to not exercise (18.5% of $121=18.5 \times 121/100=22.4$)

Sex	Exercise		No exercise		Total	
	No.	%	No.	%	No.	%
Male	33.4	81.5	7.6	18.5	41	100
Female	98.6	81.5	22.4	18.5	121	100
Total	132	81.5	30	18.5	162	100

Expected table

- Only numbers, without percentages

Sex	Exercise	No exercise	Total
Male	33.4	7.6	41
Female	98.6	22.4	121
Total	132	30	162

Quick way

Expected number = $\frac{\text{Row total} \times \text{Column total}}{\text{Overall total}}$

Observed

Sex	Exercise	No exercise	Total
Male	31	10	41
Female	101	20	121
Total	132	30	162

$$41 \times 132 / 162 = 33.4$$

Sex	Exercise	No exercise	Total
Male	33.4		41
Female			121
Total	132	30	162

Quick way

Expected number = $\frac{\text{Row total} \times \text{Column total}}{\text{Overall total}}$

Observed

Sex	Exercise	No exercise	Total
Male	31	10	41
Female	101	20	121
Total	132	30	162

$$41 \times 132 / 162 = 33.4$$

$$41 \times 30 / 162 = 7.6$$

Expected

Sex	Exercise	No exercise	Total
Male	33.4	7.6	41
Female			121
Total	132	30	162

Quick way

Expected number = Row total x Column total
Overall total

Observed

Sex	Exercise	No exercise	Total
Male	31	10	41
Female	101	20	121
Total	132	30	162

$$41 \times 132 / 162 = 33.4$$

$$41 \times 30 / 162 = 7.6$$

$$121 \times 132 / 162 = 98.6$$

Expected

Sex	Exercise	No exercise	Total
Male	33.4	7.6	41
Female	98.6		121
Total	132	30	162

Quick way

Expected number = $\frac{\text{Row total} \times \text{Column total}}{\text{Overall total}}$

Observed

Sex	Exercise	No exercise	Total	
Male	31	10	41	$41 \times 132 / 162 = 33.4$ $41 \times 30 / 162 = 7.6$
Female	101	20	121	$121 \times 132 / 162 = 98.6$
Total	132	30	162	$121 \times 30 / 162 = 22.4$

Expected

Sex	Exercise	No exercise	Total
Male	33.4	7.6	41
Female	98.6	22.4	121
Total	132	30	162

Step 2 – calculate χ^2

Compare each observed value with each expected value

	Sex	Exercise	No exercise	Total
Observed	Male	31	10	41
	Female	101	20	121
	Total	132	30	162

	Sex	Exercise	No exercise	Total
Expected	Male	33.4	7.6	41
	Female	98.6	22.4	121
	Total	132	30	162

and obtain χ^2 test statistic.

$$\chi^2 = \sum \{(O-E)^2/E\}$$

- Compare each observed value with each expected value and obtain χ^2 test statistic.
- $\chi^2 = \sum \{(O-E)^2/E\}$
- Calculate $(O-E)^2/E$ for each cell and sum over all cells
- $\chi^2 = (31 - 33.4)^2/33.4 + (10 - 7.6)^2/7.6 + (101 - 98.6)^2/98.6 + (20 - 22.4)^2/22.4 = \mathbf{1.25}$

Step 3 - Obtain p-value

- Refer χ^2 value to tables of chi-squared distribution
- Need “degrees of freedom”, v
- $v = (r - 1) \times (c - 1)$
r = no. of rows, c = no. of columns.
- In example, $r = c = 2$, so $v = (2-1) \times (2-1) = 1$
- Refer to table, $\chi^2 = 1.25$, d.f. = 1

Percentage points of the χ^2 distribution.

d.f.	P value							
	0.5	0.25	0.1	<u>0.05</u>	0.025	0.01	0.005	0.001
<u>1</u>	0.45	★ 1.32	2.71	<u>3.84</u>	5.02	6.63	7.88	10.83
2	1.39	2.77	4.61	5.99	7.38	9.21	10.60	13.82
3	2.37	4.11	6.25	7.81	9.35	11.34	12.84	16.27
4	3.36	5.39	7.78	9.49	11.14	13.28	14.86	18.47
5	4.35	6.63	9.24	11.07	12.83	15.09	16.75	20.52
6	5.35	7.84	10.64	12.59	14.45	16.81	18.55	22.46
7	6.35	9.04	12.02	14.07	16.01	18.48	20.28	24.32
8	7.34	10.22	13.36	15.51	17.53	20.09	21.96	26.13
9	8.34	11.39	14.68	16.92	19.02	21.67	23.59	27.88
10	9.34	12.55	15.99	18.31	20.48	23.21	25.19	29.59
11	10.34	13.70	17.28	19.68	21.92	24.73	26.76	31.26
12	11.34	14.85	18.55	21.03	23.34	26.22	28.30	32.91
13	12.34	15.98	19.81	22.36	24.74	27.69	29.82	34.53
14	13.34	17.12	21.06	23.68	26.12	29.14	31.32	36.12
15	14.34	18.25	22.31	25.00	27.49	30.58	32.80	37.70
16	15.34	19.37	23.54	26.30	28.85	32.00	34.27	39.25
17	16.34	20.49	24.77	27.59	30.19	33.41	35.72	40.79
18	17.34	21.60	25.99	28.87	31.53	34.81	37.16	42.31
19	18.34	22.72	27.20	30.14	32.85	36.19	38.58	43.82
20	19.34	23.83	28.41	31.41	34.17	37.57	40.00	45.32

- In example, $r = c = 2$, so $v = (2-1) \times (2-1) = 1$
- From table, χ^2 value of 1.25, $P > 0.05$

Step 4 - Interpret p-value

- No evidence of association
- Accept H_0

Quick method for χ^2

- There is a quick formula to test for association in 2x2 table
- If we label cells of 2x2 table as follows:

a b | e
 c d | f

 g h | N

Sex	Exercise	No exercise	Total
Male	31 (a)	10 (b)	41(e)
Female	101 (c)	20 (d)	121 (f)
Total	132 (g)	30 (h)	162 (N)

- Then easiest way to calculate χ^2 is using:

$$\chi^2 = \frac{(|ad - bc|)^2 \times N}{efgh} = \frac{(31 \times 20 - 101 \times 10)^2 \times 162}{41 \times 121 \times 132 \times 30} = 1.25$$

SUMMARY

What to do when confronted with categorical data?

- **6 Step Guide....**

Step 1: Construct 2-way table to display data

Step 2: Calculate row (independent) %'s

Step 3: Carry out (O-E) χ^2 test of association (or quick formula for 2x2 tables only)

Step 4: Calculate degrees of freedom for χ^2 test

Step 5: Refer to tables to obtain P-value

Step 6: Interpret p-value

Another example – Tumor response

Observed

Drug	Tumor response		Total
	Yes	No	
Drug A	49 (53%)	44	93
Drug B	18 (20%)	73	91
Total	67 (36%)	117	184

Expected

Drug	Tumor response		Total
	Yes	No	
Drug A			93
Drug B			91
Total	67 (36%)	117	184

▪

Another example – Tumor response

Observed

Drug	Tumor response		Total
	Yes	No	
Drug A	49 (53%)	44	93
Drug B	18 (20%)	73	91
Total	67 (36%)	117	184

Expected

Drug	Tumor response		Total
	Yes	No	
Drug A	33.86	59.4	93
Drug B	33.14	57.86	91
Total	67 (36%)	117	184

$$\chi^2 = (49 - 33.86)^2/33.86 + (18 - 33.14)^2/33.14 + (44 - 59.14)^2/59.14 + (73 - 57.86)^2/57.86 = 21.52.$$

Quick formula

Drug	Tumor response		Total
	Yes	No	
Drug A	49 (53%)	44	93
Drug B	18 (20%)	73	91
Total	67 (36%)	117	184

$$\chi^2 = \frac{(|ad - bc|)^2 \times N}{efgh} = \frac{(49 \times 73 - 44 \times 18)^2 \times 184}{93 \times 91 \times 67 \times 117} = 21.51$$

- χ^2 of 21.52
- $r = c = 2$, so $(2-1) \times (2-1) = 1$ d.f.
- $p < 0.05$

Percentage points of the χ^2 distribution.

<u>df.</u>	<i>P</i> value							
	0.5	0.25	0.1	<u>0.05</u>	0.025	0.01	0.005	0.001
<u>1</u>	0.45	1.32	2.71	<u>3.84</u>	5.02	6.63	7.88	10.83
2	1.39	2.77	4.61	5.99	7.38	9.21	10.60	13.82
3	2.37	4.11	6.25	7.81	9.35	11.34	12.84	16.27
4	3.36	5.39	7.78	9.49	11.14	13.28	14.86	18.47
5	4.35	6.63	9.24	11.07	12.83	15.09	16.75	20.52
6	5.35	7.84	10.64	12.59	14.45	16.81	18.55	22.46
7	6.35	9.04	12.02	14.07	16.01	18.48	20.28	24.32
8	7.34	10.22	13.36	15.51	17.53	20.09	21.96	26.13
9	8.34	11.39	14.68	16.92	19.02	21.67	23.59	27.88
10	9.34	12.55	15.99	18.31	20.48	23.21	25.19	29.59
11	10.34	13.70	17.28	19.68	21.92	24.73	26.76	31.26
12	11.34	14.85	18.55	21.03	23.34	26.22	28.30	32.91
13	12.34	15.98	19.81	22.36	24.74	27.69	29.82	34.53
14	13.34	17.12	21.06	23.68	26.12	29.14	31.32	36.12
15	14.34	18.25	22.31	25.00	27.49	30.58	32.80	37.70
16	15.34	19.37	23.54	26.30	28.85	32.00	34.27	39.25
17	16.34	20.49	24.77	27.59	30.19	33.41	35.72	40.79
18	17.34	21.60	25.99	28.87	31.53	34.81	37.16	42.31
19	18.34	22.72	27.20	30.14	32.85	36.19	38.58	43.82
20	19.34	23.83	28.41	31.41	34.17	37.57	40.00	45.32



Another example – More than 2X2

Age group	Fever after operation						Total	
	Mild		Moderate		Severe		No.	%
	No.	%	No.	%	No.	%		
<30 Y	37		14		12		63	
30-45 Y	18		17		19		54	
>45 Y	24		14		10		48	
Total	79		45		41		165	

- **Distribution of fever after operation differs according to age group**
- **but.....by more than we expect by chance??.....**

Age group	Fever after operation						Total	
	Mild		Moderate		Severe		No.	%
	No.	%	No.	%	No.	%		
<30 Y	37	59	14	22	12	19	63	100
30-45 Y	18	33	17	31	19	35	54	100
>45 Y	24	50	14	29	10	21	48	100
Total	79		45		41		165	

Observed

Age group	Fever after operation			Total
	Mild	Moderate	Sever	
<30 Y	37	14	12	63
30-45 Y	18	17	19	54
>45 Y	24	14	10	48
Total	79	45	41	165

Expected

Age group	Fever after operation			Total
	Mild	Moderate	Sever	
<30 Y	30.16	17.18	15.65	63
30-45 Y	25.85	14.73	13.42	54
>45 Y	22.98	13.09	11.93	48
Total	79	45	41	165

- **Compare each expected value with each observed value and obtain χ^2 test statistic.**
- **$\chi^2 = \Sigma \{(O-E)^2/E\}$**
- **Calculate $(O-E)^2/E$ for each cell and sum over all cells**
- **$\chi^2 = (37 - 30.16)^2/30.16 + (18 - 25.85)^2/25.85 + \dots + (10 - 11.93)^2/11.93 = 8.47$**

- **Degrees of freedom**

$$v = (r - 1) \times (c - 1) \quad r = \text{no. of rows, } c = \text{no. of columns.}$$

- **In example, $r = c = 3$, so $v = (3-1) \times (3-1) = 4$**
- **Refer to table, $\chi^2 = 8.47$, d.f. =4**
- **From table, χ^2 value of 8.47, $0.05 < P < 0.10$**
- **No evidence of association**
- **Note: We can not use quick formula here (not 2x2 table)**

Percentage points of the χ^2 distribution.

<u>df.</u>	<i>P</i> value							
	0.5	0.25	0.1	<u>0.05</u>	0.025	0.01	0.005	0.001
1	0.45	1.32	2.71	3.84	5.02	6.63	7.88	10.83
2	1.39	2.77	4.61	5.99	7.38	9.21	10.60	13.82
3	2.37	4.11	6.25	7.81	9.35	11.34	12.84	16.27
<u>4</u>	3.36	5.39	7.78	<u>9.49</u>	11.14	13.28	14.86	18.47
5	4.35	6.63	9.24	11.07	12.83	15.09	16.75	20.52
6	5.35	7.84	10.64	12.59	14.45	16.81	18.55	22.46
7	6.35	9.04	12.02	14.07	16.01	18.48	20.28	24.32
8	7.34	10.22	13.36	15.51	17.53	20.09	21.96	26.13
9	8.34	11.39	14.68	16.92	19.02	21.67	23.59	27.88
10	9.34	12.55	15.99	18.31	20.48	23.21	25.19	29.59
11	10.34	13.70	17.28	19.68	21.92	24.73	26.76	31.26
12	11.34	14.85	18.55	21.03	23.34	26.22	28.30	32.91
13	12.34	15.98	19.81	22.36	24.74	27.69	29.82	34.53
14	13.34	17.12	21.06	23.68	26.12	29.14	31.32	36.12
15	14.34	18.25	22.31	25.00	27.49	30.58	32.80	37.70
16	15.34	19.37	23.54	26.30	28.85	32.00	34.27	39.25
17	16.34	20.49	24.77	27.59	30.19	33.41	35.72	40.79
18	17.34	21.60	25.99	28.87	31.53	34.81	37.16	42.31
19	18.34	22.72	27.20	30.14	32.85	36.19	38.58	43.82
20	19.34	23.83	28.41	31.41	34.17	37.57	40.00	45.32

Questions??