



## **Practical Endocrinology**

ESR

Zhikal O. Khudur

Gasheen Bakhtiyar

Endocrinology BIO

Spring semester

Week 5



26/ 02 /2024

# Erythrocyte sedimentation rate (ESR)

- **Introduction and principle:**

➤ When blood, mixed with anticoagulant, is allowed to stand in a special, narrow vertical tube, for a period, RBCs settle down (sediment) leaving clear plasma above.

➤ **ESR is the rate at which RBCs sediment.** It is measured by the distance that the RBCs have settled down, in millimeters, in a given period, which is usually one hour.



- **Rouleaux** formation...RBCs arranged one on the top of the other....the surface area is reduced, and the weight is increased, leading to increased rapidity of sedimentation.
  - Pro-sedimentation factors, mainly **fibrinogen** and **globulins**
  - Factors resisting sedimentation, namely the **negative charge** of the erythrocytes
  - **The balance between above factors governs the ESR.**
- **Therefore, the factors that increase rouleaux formation increase the ESR, and those factors that decrease rouleaux formation decrease the ESR.**

# Factors which affect ESR

## 1. Plasma proteins

- Increased concentrations of **fibrinogen** or **globulin** & **acute phase proteins** (increased in acute inflammation or tissue injury).....all increase aggregating ability, therefore, increase rouleaux formation and **increase** ESR.
- Increased concentrations of **albumin** in plasma increase RBC negative charge so reduce rouleaux formation and **decrease** ESR.

**2. Concentration of RBCs:** Increased RBC concentration or high Hct increases the **viscosity** of blood thus increasing the **resistance** to sedimentation, which leads to **decrease in ESR** and vice versa.

- **Viscosity** is important in circulatory function because it partially governs the **flow** of blood through the vessels.
- Increased **viscosity**.....increased resistance....decrease **blood flow** and vice versa.
- Either of these conditions puts a strain on the heart that may lead to serious cardiovascular problems if not corrected.

**3. Shape of RBCs:** RBCs with abnormal or irregular shapes such as sickle cells or spherocytes interfere with rouleaux formation leading to decreased ESR .

- **ESR is not a specific test** so it doesn't give an indication about the **type** or **site** of disease but it's used for the **follow up** of the patient and the prognosis of the condition.
- Affected by other conditions besides inflammation thus used in **conjunction** with other tests.
- In people **above 60** years of age, higher values are not necessarily abnormal.

# ESR method, material and instruments

- **Westergren pipette** it is glass tube (30 cm in length & 2.55 mm in width), opened at both ends & graduated from **0-200 mm**.
- Westergren pipette rack equipped with leveling screws
- 3.8% Sodium citrate as an anticoagulant in a **ratio of 1:4** with blood
- Syringe for the withdrawal of blood from the vein of the patient.

# Procedure:-

- Withdraw **2 cc of blood** from the patient's vein using the syringe.
- Put **0.4 cc** of **sodium citrate** in a plain test tube.
- Immediately add **1.6 ml blood** from the syringe to the plain tube making a dilution of **1:4** ratio and shake it for mixing.
- Fill the Westergren pipette to exactly the **0 mark** making certain that there are **no air bubbles** at all in the blood
- Place the pipette vertically on the rack and leave it to stand still undisturbed for **one hour**.
- At the end of the 60 minutes read the **no. of millimeters the RBC's have fallen** (i.e. the height of clear plasma above the upper limit of the column of sedimentary cells) the result is the ESR in **mm/ in 1 hour**.





# Technical errors

- A **tilted** ESR tube.
- Increase **temperature**.
- Sample **dilution**.
- **Inadequate** anticoagulation with clotting of the blood sample (consequently will consume fibrinogen).....result will?

# Medical application

## The **ESR increase** in:

- **Physiological**
  - pregnancy, menstruation & parturition
- **Pathological**
  - anemia, infections, malignancy, and inflammation.

## The **ESR decrease** in:

- polycythemia, spherocytosis, and sickle cell anemia.

## Normal range for ESR test

Men under 50 years old: less than 15 mm/hr

Men over 50 years old: less than 20 mm/hr.

Women under 50 years old: less than 20 mm/hr.

Women over 50 years old: less than 30 mm/hr.