

Microbial growth and metabolism

Prof.Dr.Sewgil S.Anwer

Lec.4-5

2024-2025


Microbial growth and microbial growth requirement

Microbial growth

Microbial growth is the increase in number of cells, not cell size.

Growth in colonies

- A pure culture contains only one species or strain.
- A colony is a population of cells arising from a single cell or spore or from a group of attached cells.
- a unit used to estimate the number of viable **microbial** cells is colony forming unit (CFU)



Bacterial Division

The normal reproductive method of bacteria is **binary fission**, in which a single cell divides into two identical cells

Bacterial Division


The normal reproductive method of bacteria is binary fission, in which a single cell divides into two identical cells.

1 Cell elongates and DNA is replicated

2 Cell wall and plasma membrane begin to grow inward

3 Cross-wall forms completely around divided DNA

4 Cells separate



(a) A diagram of the sequence of cell division.

Growth rate

The number of generation per hour.

Generation Time


The **time required** for a cell to divide or a **population to double** is known as the generation time.

<i>Escherichia coli</i>	10 minutes
<i>Mycobacterium tuberculosis</i>	15 hours

The generation time depend upon:

- 1- The nutrient in the medium.
- 2- Physical condition (pH, temp. etc.)


Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.



(a)

Q/- what is the generation number of bacteria when the number of cell is 8


- If the generation number of bacteria was 4 what is the Exponential value?



Mathematical Expression of Growth

Number of Generations

Cell Numbers



Bacterial growth curve

- All microorganisms undergo similar growth patterns

Each growth curve has 4 phases:

1- Lag phase

2- Log phase

3- Stationary phase

4- Death or decline phase

Between each phases there is a **transitional phase** is represent the time require by all the cell before get to inter the new phase.

1- Lag phase:

- The number of the population remains constant.
- Microorganism start to adapted itself to the environment.
- The lag phase is generally longer if the cells are taken from an old or refrigerated culture.
- If the cells are taken from young, new growing culture (microbial population) and inoculated to a fresh medium the lag phase may be short or even absent

2- Log phase (logarithmic phase or exponential phase):

- The bacteria multiply at the fastest rate possible under the conditions provided.
- Most research is performed on cells during log phase
- This phase is called log phase because the logarithm of the bacterial mass increases linearly with time,
- and exponential growth phase because the number of cells increases as an exponential function of $2n$ (i.e. 2^1 , 2^2 , 2^3 , 2^4 , 2^5 and so on).
- The portion of the growth curve where rapid growth of bacteria is observed is known as **Logarithmic phase**.

3- Stationary phase:

- Growth levels off.
- Cells per volume does not increase or decrease.
- Growth rate = Death rate.

Due to


- Depletion of nutrients
- Increase in waste products

4- Death phase:

- The number of deaths exceeds the number of new cells formed
- Cells per volume decreases
- Due to
 - Very low concentrations of nutrients
 - Very high concentrations of waste products

Standard Growth Curve

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.



Nutritional Requirements for Microorganisms:

-Water (preservation of a microbial culture from drying)

-Energy:


Phototroph- Energy from sun light (cyanobacteria)

Chemotroph- Energy from chemicals (Nitrobacter, **Sulphur bacteria**)

Heterotroph- from carbohydrate and other organic material

-Carbon:

- Autotrophs- carbon from carbon dioxide- (inorganic carbon)
- Heterotroph- carbon from organic carbon
 - e.g. Carbohydrates, lipids, protein.



- Essential elements

- Hydrogen (H), Sulfur (S), Oxygen (O), Phosphorous (P)
- Nitrogen (N)
 - commonly supplied as ammonia (NH₄)
 - some microbes fix atmospheric nitrogen (N₂)

- Trace elements

- Required in small amounts
- Copper (Cu), Zinc (Zn), Selenium (Se)

Environmental requirements for growth

- Temperature, pH, Oxygen, Carbon dioxide, Osmotic pressure, Hydrostatic pressure

- **Temperature**

- Psychrophiles – less than 20 °C
- Mesophiles – 20 - 45 °C
- Thermophiles – 45 – 80 °C
- Extreme thermophiles – more than 85 °C

- **Psychrotroph- 0-40 (20) °C**

- **pH** (- log [H⁺])

Low pH = acid, High pH = basic or alkaline

- Acidophiles - below pH 5.5 (Acidobacteriota, Mucor)
- Neutrophiles – at pH 6 – 8
- Alkalophiles – above pH 8 (Thiohalospira alkaliphila)

- **Molecular oxygen**

Microbe vary greatly in sensitivity to oxygen.

- Aerobes – microbes which require oxygen.
- Facultative anaerobes – microbes which can grow in presence or absence of oxygen.
- Obligate Anaerobes – which do not utilize oxygen and are killed by oxygen.
- Aerotolerant anaerobe is an organism that tolerates the presence of oxygen but does not require it for growth.
- Microaerophiles – required 3 – 15 % oxygen.

Chemical Requirements for Growth: **Oxygen**

O₂ requirements vary greatly




Table 6.1: The Effects of Oxygen on the Growth of Various Types of Bacteria

- **Carbon dioxide**


- Capnophiles : 3 – 10 % carbon dioxide
- Many microaerophiles are also capnophiles




Osmotic Pressure

High osmotic pressure (hypertonic) removes water causing plasmolysis – inhibits growth i.e. salt as preservative

Low osmotic pressures (hypotonic) cause water to enter and can cause lysis



(d) Hypotonic (hypoosmotic) solution—water moves into the cell and may cause the cell to burst if the wall is weak or damaged (osmotic lysis)



(e) Hypertonic (hyperosmotic) solution—water moves out of the cell, causing its cytoplasm to shrink (plasmolysis)

Quantitative methods for measuring growth of bacteria:

The growth of bacteria can be determined by numerous techniques based on one or more of the following types of measurement:

1- Cell count

- a- microscopy or by using electronic particle counter.
- b- colony count or number (plate count method, MPN)

2- Cell mass

- a- weighting (dry weight)
- b- Measurement of cell nitrogen
- c- Indirectly by turbidity with culture , directly without culture

3- Cell activity- indirectly by relating the degree of biochemical activity to the size of population Such as measurement of utilizing O₂

Direct methods :


With direct methods we count individual cells or colonies that are assumed to have apart or arise in through the division of a single cell.

1- Counting Chamber (Hemocytometer) :


The hemocytometer is a specialized microscope slide used to count cells.

The center portion of the slide has etched grids (H) with precisely spaced lines.







Neubauer's Chamber or Haemocytometer



CORNER SQUARE (ENLARGEMENT)



Howard
Cell



Count cells on top and left touching middle line (O).
Do not count cells touching middle line at bottom and right (Ø).

Steps of Counting of cell using hemocytometer

1- percentage of viable cell= **Total viable cell/ total cell *100**

2-Average of cell per square = **total viable/ average**

3- Dilution factor =**Final volume / volume of cell**

4- Concentration (viable cells/ ml)=**Average of cell* dilution factor***

10000 (10⁴)

Suppose you have 200 ml of sample diluted with 200 ml stain. you calculated the number of viable cell as 36 and the number of a viable cell (dead cell) was 6 and the average of cell per square was 6.

Answer the following questions

1. Calculate Percentage of viable cell
2. Find the concentration of viable cell/ml

Answer

$200+200= 400$ total volume

D. f= final volume/ volume .of cell($400/200=2$)

Total cell= $36+6=42$

Percentage of viable cell= total viable cell/ total cell*100
 $= 36/42*100= 85.7$

concentration of viable cell/ml= Av. Of cell * dilution factor* 10^4
 $=6*2* 10^4 = 120,000$

2- Coulter Counter :

electronic counting (this machine detects the difference in current as individual microorganisms pass through a small orifice).

It is Very fast , easy to use but;

Very EXPENSIVE.




3- Viable count assays (Colony Counting) :

Colony counting after plating dilutions of the sample onto growth medium.

Standard plate counts using spread and pour plate techniques (cfu for “colony forming unit”).

two viable count assays:

1 - Spread plates 2- Pour plates



(a) The pour plate method

(b) The spread plate method

1 Inoculate empty plate

1.0 or 0.1 ml

0.1 ml

Bacterial dilution

1 Inoculate plate containing solid medium

2 Add melted nutrient agar

2 Spread inoculum over surface evenly

3 Swirl to mix

4 Colonies grow in and on solidified medium


3 Colonies grow only on surface of medium



Indirect Method :

Indirect methods often rely on the results of metabolic tests or other growth characteristics. And it's to:

- Measurement of metabolic activity.
- Gas or Acid Production.



- **Turbidity using a spectrophotometer.**

spectrophotometry, using a spectrophotometer .

These Indirect counts depend on:

- The effects of the organisms to estimate their numbers.
- As organisms grow they make the nutrient broth turbid.
- This turbidity can be measured with a colorimeter

